

## **Ways to Support Yourself During Social Distancing, Quarantine, and Isolation**

*(Sourced from SAMHSA publication, US Department of Health and Human Services)*

### **UNDERSTAND THE RISK**

Consider the real risk of harm to yourself and others around you. The public perception of risk during a situation such as an infectious disease outbreak is often inaccurate. Media coverage may create the impression that people are in immediate danger when really the risk for infection may be very low. Take steps to get the facts:

- Stay up to date on what is happening, while limiting your media exposure. Avoid watching or listening to news reports 24/7 since this tends to increase anxiety and worry. Remember that children are especially affected by what they hear and see on television.
- Look to credible sources for information on the infectious disease outbreak (see page 2 for sources of reliable outbreak-related information).

### **BE YOUR OWN ADVOCATE**

Speaking out about your needs is particularly important if you are in quarantine, since you may not be in a hospital or other facility where your basic needs are met. Ensure you have what you need to feel safe, secure, and comfortable.

- Work with local, state, or national health officials to find out how you can arrange for groceries and toiletries to be delivered to your home as needed.
- Inform health care providers or health authorities of any needed medications and work with them to ensure that you continue to receive those medications.

### **EDUCATE YOURSELF**

Health care providers and health authorities should provide information on the disease, its diagnosis, and treatment.

- Do not be afraid to ask questions—clear communication with a health care provider may help reduce any distress associated with social distancing, quarantine, or isolation.
- Ask for written information when available.

- Ask a family member or friend to obtain information in the event that you are unable to secure this information on your own.

### **CONNECT WITH OTHERS**

Reaching out to people you trust is one of the best ways to reduce anxiety, depression, loneliness, and boredom during social distancing, quarantine, and isolation. You can:

- Use the telephone, email, text messaging, and social media to connect with friends, family, and others.
- Talk “face to face” with friends and loved ones using Skype or FaceTime.
- If approved by health authorities and your health care providers, arrange for your friends and loved ones to bring you newspapers, movies, and books.
- Sign up for emergency alerts via text or email to ensure you get updates as soon as they are available.
- Use the Internet, radio, and television to keep up with local, national, and world events. Sources for reliable outbreak-related information include Centers for Disease Control and Prevention (<http://www.cdc.gov> ) and World Health Organization (<http://www.who.int/en> )

### **USE PRACTICAL WAYS TO COPE AND RELAX**

- Relax your body often by doing things that work for you—take deep breaths, stretch, meditate or pray, or engage in activities you enjoy.
- Pace yourself between stressful activities, and do something fun after a hard task.
- Talk about your experiences and feelings to loved ones and friends, if you find it helpful.
- Maintain a sense of hope and positive thinking; consider keeping a journal where you write down things you are grateful for or that are going well.

### **AFTER SOCIAL DISTANCING, QUARANTINE, OR ISOLATION**

You may experience mixed emotions, including a sense of relief. If you were isolated because you had the illness, you may feel sadness or anger because friends and loved ones may have

unfounded fears of contracting the disease from contact with you, even though you have been determined not to be contagious.

The best way to end this common fear is to learn about the disease and the actual risk to others. Sharing this information will often calm fears in others and allow you to reconnect with them.

If you or your loved ones experience symptoms of extreme stress—such as trouble sleeping, problems with eating too much or too little, inability to carry out routine daily activities, or using drugs or alcohol to cope—speak to a health care provider.